

## TREATMENT

# Options

Cysts that cause no symptoms and appear as “simple” cysts on imaging require no treatment. Patients can be reassured that they are not cancerous, nor does the presence of cysts increase the risk of cancer.


Painful, large simple cysts may be aspirated for comfort. This involves draining the cyst with a fine needle, usually under ultrasound guidance (fine needle aspiration or FNA). The fluid obtained is commonly watery and strawcoloured. Occasionally, cysts may recur after FNA or new cysts can develop in the breast tissue nearby.

Complex cysts need to be accurately biopsied, usually by means of a core biopsy to obtain sufficient tissue for analysis. The cyst is then managed according to the pathology result. Surgery is not normally recommended for managing simple cysts, however it may be required for complex cysts if the pathology result suggests a cancerous change or if the cyst cannot be accurately sampled by needle biopsy.

It is important that women, particularly those who often get cysts, do not presume that a new lump is just another cyst. Medical attention should always be sought for any new breast lump.



breast cancer  
research centre-wa  
Incorporating Perth Breast Cancer Institute



### Perth Breast Cancer Institute

Hollywood Consulting  
Centre Suite 404-405, Level 4  
91 Monash Avenue, Nedlands 6009

 (08) 6500 5576  [reception@bcrc-wa.com.au](mailto:reception@bcrc-wa.com.au)

Published by Breast Cancer Research Centre-WA August 2021



breast cancer  
research centre-wa  
Incorporating Perth Breast Cancer Institute

## BREAST

# Cysts

## WHAT ARE

# Breast Cysts?


Breast cysts are fluid-filled sacs inside the breast tissue. They are very common, occurring in up to 50% of women in the 30-50 year age group. Cysts are affected by hormonal changes and therefore they are most common in women during the reproductive years. Cysts often disappear after the menopause but they may be seen in older women taking menopause hormonal therapy.

Cysts form due to an imbalance between fluid production and absorption in the small breast ducts and lobules. They may be single or multiple and occur in one or both breasts. They may also fluctuate in size and number over time. Larger cysts may be felt by a woman as a rounded, smooth, mobile lump. Smaller cysts often cause no symptoms at all and may be found incidentally on breast ultrasound or mammogram. Fortunately, most cysts are not cancerous.

## WHEN TO SEE A

# Doctor

Women usually seek medical attention because of a lump or its associated pain or discomfort. Rarely, cysts may become inflamed resulting in increased pain and redness of the overlying skin. It is not possible for physical examination alone to accurately distinguish between a benign cyst and other types of breast lumps so it is always important for a woman to promptly report any new breast symptom to her doctor.



*Ultrasound image showing a breast cyst with well-defined margins.*

## DIAGNOSIS

Diagnosis of breast cysts entails a clinical breast examination and breast imaging, usually in the form of an ultrasound and/or mammogram. The choice of initial imaging modality depends on the women's age and other factors such as pregnancy or breastfeeding. On mammogram, cysts appear as smooth rounded lumps. On ultrasound, cysts may be described as "simple", containing only fluid within a thin-walled sac or "complex". Complex cysts may have other features such as a thickened wall or solid components in addition to fluid. These are usually biopsied to exclude the small chance of breast cancer.

